



Robotic assisted sleeve lobectomy with the use of barbed sutures

Jan-Hendrik Egberts, Thorben Möller, Charlotte Hauser, Jan-Henrik Beckmann, Mark Schlemminger, Thomas Becker

Department of General, Visceral-, Thoracic-, Transplantation-, and Pediatric Surgery, Kurt-Semm Center for Laparoscopic and Robotic Assisted Surgery, University Hospital Schleswig Holstein, Campus Kiel, Kiel, Germany

Correspondence to: Jan-Hendrik Egberts, MD. Department for General, Visceral-, Thoracic-, Transplantation-, and Pediatric Surgery; University Hospital Schleswig Holstein, Campus Kiel, Arnold Heller Str. 3, 24105 Kiel, Germany. Email: jan-hendrik.egberts@uksh.de.

Abstract: Anastomosis in bronchoplasty using continuous suturing has become widely accepted. However, especially in VATS and robotic assisted thoracic surgery (RATS) it is difficult to maintain a continuous traction of suture over the circumference. In addition, the lack of haptic feedback on the current robotic systems might even further complicate the precise and constant maintenance of traction. Recent reports of using absorbable barbed suture materials in open and VATS surgery were published with promising results. We adopted and standardized the application of barbed sutures (Stratafix, Johnson & Johnson, USA). There were no intraoperative difficulties such as cutting, loosening or tailing the suture.

Keywords: Robotic sleeve lobectomy; bronchoplasty; absorbable barbed suture; robotic assisted thoracic surgery (RATS)

Received: 19 April 2018; Accepted: 08 May 2018; Published: 04 June 2018.

doi: 10.21037/jovs.2018.05.19

View this article at: <http://dx.doi.org/10.21037/jovs.2018.05.19>

Introduction

In bronchoplastic operations of the lung, the reconstruction of the airway is usually performed by applying an anastomosis of the bronchus. This is traditionally created in interrupted sutures that are considered to be precise, safe, and reliable (1). However, the placement of interrupted sutures can be complex and time consuming. Usually knots of the back wall of the anastomosis are placed intraluminal whereas the front wall is knotted outside. But especially the intraluminally placed nodes of the inner wall might also lead to a stenosis of the lumen with consecutive accumulation of the secretion (2).

Bronchial anastomosis using continuous suturing has recently become widely accepted, especially in VATS and robotic assisted thoracic surgery (RATS) (3,4). However, the difficulty is to maintain a continuous traction of suture. This is even more difficult in the rigid tissue of the bronchus cartilage. In addition, the lack of haptic feedback on the current robotic systems might further complicate the precise and continuous maintenance of traction or even lead to a bunk of the suture.

Recent reports have demonstrated the usefulness of

so-called absorbable barbed sutures in laparoscopic and thoracoscopic surgery (5-7). It is a novel resorbable suture that was originally developed for dermal closure and has a unique barbed structure that prevents loosening of the suture. Based on our own positive experience with this suture in robotic esophageal and bariatric surgery, we have also used this suture in bronchoplastic surgery (8). We report for the first time our clinical experience of robotic assisted sleeve lobectomy with bronchoplasty using barbed continuous sutures.

Surgical technique

The oncological resection is performed in a robotic assisted approach (RATS) as previously described using the DaVinci Si or Xi System (Intuitive Surgery, Sunnyvale, CA, USA) (9). We apply the four-arm technique, where the trocars are placed in the 8th or 9th intercostal space. Systematic hilar and mediastinal lymph node dissection is performed until bronchial reconstruction commenced. After transection of the proximal and distal bronchus, bronchial anastomosis is performed as follows (sample for right upper lobe sleeve resection): we use the large robotic needle driver and




Figure 1 Reconstruction of the airway: right main bronchus and right intermediate bronchus. (A) Start of the anastomosis: outside to inside directed stich at the right intermediate bronchus; (B) inside to outside directed stich at right main bronchus; (C) catching the looping of the suture.




Figure 2 (A) Suturing the back wall (pars membranacea); (B) the anastomosis is semicircular continued through the cartilaginous part).




Figure 3 Suture of the front side of the airway. (A) Start with the second suture; (B) suturing the front side; (C) “Z”-Stich for fixation

Cadiere Forceps (Intuitive Surgery, Sunnyvale, CA, USA). For the anastomosis we prefer a monofilament barbed suture (Stratafix 3/0, Ethicon Endosurgery, OH, USA). The first suture is placed on the posterior part of the right bronchus intermedius from out to in to place the knot outside the airway. The corresponding stitch is performed on the right main stem bronchus from in to out, this places the knot in the cartilage of the main stem bronchus close to the membranous part (*Figure 1*). It is important to perform the first stitch in the area of the cartilage to ensure adequate suture retention. The anastomosis is now continued through

the cartilaginous part, while slowly bringing the two cut ends together (*Figure 2*). When half of the circumference is completed, a second suture is used (*Figure 3*). This suture begins at the cranial dorsal part of the cartilage to the front side until the first suture is reached. An additional stitch is then placed and locked in the membrane—cartilage junction in the same fashion. Since we did not knot the barbed suture due to its rigidity we placed a single “Z”-Stich with Vicryl 3/0 at the end of bot barbed sutures in order to fixate them in position. The incongruity of both lumens can be well compensated by different seam distances on both sides.

We do not telescope the anastomosis. We perform a leak test under water and cover the anastomosis with either a mediastinal tissue flap or an intercostal muscle flap. Finally, the insertion of a 24 Ch thorax drainages takes place.

Discussion

Sleeve lobectomy is one of the most difficult operations in lung surgery. Mostly this operation is done as an open procedure, only a limited number of medical centers are able to perform VATS sleeve lobectomy. The main reasons are that airway reconstruction with conventional VATS instruments is difficult and inconvenient due to the limited view and particularly a lack of freedom of movement—which is essential for sewing and knots. Therefore, airway reconstructions are often combined with mini-thoracotomies (1,2). The use of a robot outweighs these disadvantages with the enlarged three-dimensional view and angulation of the robot arm (4,10). However, currently available systems also have a significant drawback, such as the lack of haptic feedback. This might be of great importance while sewing fine structures in which continuous tension is required without tearing the suture.

In the past the bronchial anastomosis was sewed in single stitch technique. Nowadays bronchial anastomosis is increasingly performed in continuous running suture technique (1,2). The suture material used is usually a monofilament material. Because of the flexibility of these sutures, it is important to maintain continuous tension to avoid leakages. This condition is sometimes limited in the minimally invasive technology. Especially in the RATS technique, with the lack of haptic feedback, there might be the danger that these tensile forces will not be kept constant or too strong and therefore lead to a damage of the tissue or the suture.

The barbed suture Stratafix™ is a knotless tissue control device that was originally developed for soft-tissue approximation (Ethicon, Cincinnati, OH, USA). Anchors are cut into the core of the suture at 2 mm intervals, pointing away from the suture direction. Because each anchor functions to grip the sutured tissue, a stable maintenance of continuous sutures is possible. The suture thread is absorbable but maintains its tensile strength for 6 weeks (11,12). Compared to other spine stitches, the Stratafix™ thread has slightly finer and spiky barbs while, for example, the V-Loc™ (Covidien, New Haven, CT, USA) barbs appear bristling and more rigid. This is due to the manufacturing process, so the Stratafix thread has so-

called single angle cut whereas the V-Loc is constructed by dual angle cuts (13).

There are reports of the successful use of the V-Loc in the open or VATS technique for the tracheobronchial reconstruction (7). However, there are no clinical reports of the use of continuous sutures with neither the Stratafix™ suture device in tracheobronchial reconstruction so far, nor any reports of its application with the robot.

We are of the opinion that in fine and delicate tissue, as is the case with the suture of pars membranacea, preferably atraumatic sutures are mandatory. The slightly finer barbs of the Stratafix™ suture give these conditions. However, we would like to point out that it is a purely personal assessment and there are no clinical studies in this regard so far.

The overlapped suture for one or two stitches at the end is important for preventing an anastomosis detachment if the end of the suture is slipped off after surgery. We think it makes sense to place an additional stitch from the endpoint that the previous suture is trimmed and appropriately conserved. There are no data regarding the risks of biological reaction or bacterial infection, which could occur more frequently because of the device's barbed structure. Therefore, a number of additional clinical experiences are needed to confirm its safety in bronchial anastomosis.

Acknowledgements

None.

Footnote

Conflicts of Interest: Jan-Hendrik Egberts is paid by Intuitive Surgery™ for proctoring surgical teams. Thomas Becker received a research grant by Intuitive Surgery™. The other authors have no conflicts of interest to declare.

Informed Consent: Written informed consent was obtained from patient for publication of this report and any associated images and videos.

References

1. Maurizi G, D'Andrilli A, Venuta F, et al. Reconstruction of the bronchus and pulmonary artery. *J Thorac Dis* 2016;8:S168-80.
2. Yu JA, Weyant MJ. Techniques of bronchial sleeve resection. *Semin Cardiothorac Vasc Anesth* 2012;16:196-202.
3. Yang R, Shao F, Cao H, et al. Bronchial anastomosis using

- complete continuous suture in video-assisted thoracic surgery sleeve lobectomy. *J Thorac Dis* 2013;5 Suppl 3:S321-2.
4. Cerfolio RJ. Robotic sleeve lobectomy: technical details and early results. *J Thorac Dis* 2016;8:S223-6.
 5. Facy O, De Blasi V, Goergen M, et al. Laparoscopic gastrointestinal anastomoses using knotless barbed sutures are safe and reproducible: a single-center experience with 201 patients. *Surg Endosc* 2013;27:3841-5.
 6. Haga N, Kurita N, Yanagida T, et al. Effects of barbed suture during robot-assisted radical prostatectomy on postoperative tissue damage and longitudinal changes in lower urinary tract outcome. *Surg Endosc* 2018;32:145-53.
 7. Nakagawa T, Chiba N, Ueda Y, et al. Clinical experience of sleeve lobectomy with bronchoplasty using a continuous absorbable barbed suture. *Gen Thorac Cardiovasc Surg* 2015;63:640-3.
 8. Egberts JH, Stein H, Aselmann H, et al. Fully robotic da Vinci Ivor-Lewis esophagectomy in four-arm technique-problems and solutions. *Dis Esophagus* 2017;30:1-9.
 9. Egberts JH, Schlemminger M, Schafmayer C, et al. Robot-assisted minimally invasive lobectomy with systematic lymphadenectomy for lung cancer. *Zentralbl Chir* 2015;140:15-6.
 10. Jo MS, Kim DY, Jeong JY, et al. Robotic sleeve lobectomy with four arms for lung cancer centrally located in the right lower lobe: a case report. *J Cardiothorac Surg* 2017;12:108.
 11. Jordan MC, Holscher-Doht S, Jakubietz MG, et al. Suture material for flexor tendon repair: 3-0 V-Loc versus 3-0 Stratafix in a biomechanical comparison ex vivo. *J Orthop Surg Res* 2014;9:72.
 12. Lin Y, Lai S, Huang J, et al. The Efficacy and Safety of Knotless Barbed Sutures in the Surgical Field: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Sci Rep* 2016;6:23425.
 13. Greenberg JA, Goldman RH. Barbed suture: a review of the technology and clinical uses in obstetrics and gynecology. *Rev Obstet Gynecol* 2013;6:107-15.

doi: 10.21037/jovs.2018.05.19

Cite this article as: Egberts JH, Möller T, Hauser C, Beckmann JH, Schlemminger M, Becker T. Robotic assisted sleeve lobectomy with the use of barbed sutures. *J Vis Surg* 2018;4:117.