



Port-access thoracoscopic anatomical left S3 segmentectomy

Hiroyuki Oizumi, Hirohisa Kato, Jun Suzuki, Akira Hamada, Kenta Nakahashi, Mitsuaki Sadahiro

Department of Surgery II, Yamagata University, Yamagata, Japan

Correspondence to: Hiroyuki Oizumi, MD, PhD. Department of Surgery II, Yamagata University, 2-2-2 Iida-Nishi, Yamagata 990-9585, Japan.

Email: hohizumi@med.id.yamagata-u.ac.jp.

Abstract: We present our method for video-assisted thoracic surgery (VATS) segmentectomy based on three-dimensional (3D) computed tomography (CT) surgical simulation especially about the left S3 segmentectomy. A 71-year-old woman was referred to our hospital for her ground-glass opacity nodule (GGN) that increased in size up to 1.0 cm in the S3 of her left lung during the postoperative follow-up after S9,10 thoracoscopic segmentectomy for right lung lower lobe adenocarcinoma in our hospital 10 years ago. As the nodule was deeply located thoracoscopic anatomical left S3 segmentectomy was performed. The operative time was 152 min, and the blood loss was 4 mL. Air leakage was not observed and the chest tube was removed on postoperative day 1. The final diagnosis was a lepidic adenocarcinoma, and she lives well without recurrence in this 43-month period after the surgery. The surgical procedure is presented.

Keywords: Lung cancer; segmentectomy; thoracoscopic

Received: 25 April 2018; Accepted: 21 June 2018; Published: 10 July 2018.

doi: 10.21037/jovs.2018.06.12

View this article at: <http://dx.doi.org/10.21037/jovs.2018.06.12>

Introduction

The indications for video-assisted thoracic surgery (VATS) segmentectomy are growing with the significant advancements in surgical techniques. However, this procedure seems to be limited for anatomically simple and easy procedures such as left upper division, lingular (S4+5) and S6 segmentectomies. Intersegmental dissection in such segmentectomies is not procedurally difficult, because the dissection can simply be performed by using stapling devices along the single intersegmental plane.

We present the surgical procedure for VATS anatomical left S3 segmentectomy according to the simulation of surgical margin and bronchovascular branching using three-dimensional (3D) computed tomography (CT) reconstruction.

Procedure

Resection of left S3 was planned as the nodule was located between the subsegmental bronchi. The bronchovascular branching pattern was identified under precise assessment of 1-mm sliced axial images and 3D reconstruction of multidetector CT data (*Figure 1*). Standard anesthesia

for thoracic surgery was performed with a double lumen endotracheal tube and general anesthesia. The patient was placed in a lateral position with the left upper extremity fixed onto the hand shelf. The surgeon stood in an anterior position to the patient.

Two flexible 5-mm ports and one 20-mm soft silicone port were inserted, and a 30° 10-mm rigid scope was used. We first incised the lung parenchyma along the V1+2 that runs between S1+2 and S3, then the V3b that runs between S3 and S4. The hilum of S3 was exposed and we isolated and divided the vein V3c that runs into the affected segment. In this patient, as the first branch of left pulmonary artery was composed of anterior artery (A3) and lingular artery (A4+5) the A3 was isolated from the A4+5 branch and ligation and division were done. The anterior segmental bronchus (B3) was looped using a monofilament polypropylene thread. A slip knot was made using the thread outside her thorax, and bilateral pure oxygen lung ventilation was performed. During the lung ventilation, the end of the thread was pulled, and the knots slipped to reach the bronchus without a knot pusher so as to trap the segmental air of the S3. The unilateral ventilation was resumed and the B3 was divided using a stapler, then dissection of hilar parenchyma was done dividing venous




Figure 1 The tumor and bronchovascular tree using 3D reconstruction of multi-detector CT data. (A) 3D-CT image. Images of arteries (red), veins (blue), bronchi (green), and tumor (magenta) are reconstructed by surgeons; (B) rotated 3D-CT image. The tumor is located between subsegmental bronchi, B3a, B3b and B3c. Black arrow, V3 (vein between S3 and S4); white arrow, V1+2 (vein between S1+2 and S3); black arrowhead, V4+5 (lingular vein). PA, pulmonary artery; PV, pulmonary vein; 3D-CT, three-dimensional computed tomography.



Figure 2 Surgical procedure of thoracoscopic left S3 segmentectomy (1).

Available online: <http://www.asvide.com/article/view/25743>

branches running into the segments. During this procedure, the inflation–deflation lines between the segments gradually become apparent. We further incised the lung parenchyma along the inflation–deflation line and finally the peripheral lung was divided using staplers. Fibrin glue was sprayed at the remnant cut surface (*Figure 2*) (1). The operative time was 152 min. The estimated blood loss was 4 mL. Air leakage was not observed, and the chest tube was removed at the first postoperative day. The final diagnosis was a lepidic adenocarcinoma. She lives well without recurrence 42 months after the left lung operation.

Comments

We previously reported the thoracoscopic anatomical

segmentectomy under the use of 3D reconstruction of multidetector row CT of the affected segments and so far, performed 335 procedures including 10 right and 7 left S3 segmentectomies. To perform the anatomical segmentectomy procedure, precise understanding of hilar bronchovascular anatomy is important under the use of multidetector CT data (2,3). As for the left S3 segmentectomy, we first catch the anatomy of V3 as the branches usually run between left upper division and lingular segment, then put weight on the running of V1+2 that runs between apical posterior segment (S1+2) and S3 under precise assessment of 1-mm sliced axial images as well as the 3D reconstruction of multidetector CT data. Anterior artery (A3) and bronchus (B3) are usually identifiable between these two veins. Thus, during the S3 segmentectomy, we first identify these veins at the hilum and dissect the lung parenchyma towards peripheral using energy sealing devices such as ultrasonically activated device or along the intersegmental veins and isolate A3 and B3.

It is beneficial to maintain the affected segment inflated for the resection of segment with multiple dissection surfaces. The merit of creating an inflation–deflation line is that the demarcation line does not disappear once it becomes visible. The jet ventilation method is developed for this purpose in thoracoscopic surgery or small thoracotomy (4). However, this method needs an extra equipment and maneuver of a bronchoscope. We developed the slip-knot method that enabled the creation of an inflation–deflation line even in anatomical subsegmental resection of which the bronchial diameter is small (5,6). Although some other

methods of visualizing the intersegmental plane (i.e., injection of dye into the target segmental bronchus using a needle or infrared visualization with intravenous injection of indocyanine green) have been reported, some processes may need special equipment or bronchoscopy skill of the anesthetist.

In conclusion, anatomical segmentectomy can be performed under precise recognition of intersegmental veins and creation of inflation-deflation line as intersegmental borders.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Informed Consent: Written informed consent was obtained from the patient for publication of this manuscript and any accompanying images.

doi: 10.21037/jovs.2018.06.12

Cite this article as: Oizumi H, Kato H, Suzuki J, Hamada A, Nakahashi K, Sadahiro M. Port-access thoracoscopic anatomical left S3 segmentectomy. *J Vis Surg* 2018;4:139.

References

- 1 Oizumi H, Kato H, Suzuki J, et al. Surgical procedure of thoracoscopic left S3 segmentectomy. *Asvide* 2018;5:604. Available online: <http://www.asvide.com/article/view/25743>
- 2 Oizumi H, Kanauchi N, Kato H, et al. Total thoracoscopic pulmonary segmentectomy. *Eur J Cardiothorac Surg* 2009;36:374-7; discussion 377.
- 3 Oizumi H, Kanauchi N, Kato H, et al. Anatomic thoracoscopic pulmonary segmentectomy under 3-dimensional multidetector computed tomography simulation: a report of 52 consecutive cases. *J Thorac Cardiovasc Surg* 2011;141:678-82.
- 4 Okada M, Mimura T, Ikegaki J, et al. A novel video-assisted anatomic segmentectomy technique: selective segmental inflation via bronchofiberoptic jet followed by cautery cutting. *J Thorac Cardiovasc Surg* 2007;133:753-8.
- 5 Oizumi H, Kato H, Endoh M, et al. Slip knot bronchial ligation method for thoracoscopic lung segmentectomy. *Ann Thorac Surg* 2014;97:1456-8.
- 6 Endoh M, Oizumi H, Kato H, et al. How to demarcate intersegmental plane with resected-segments inflation method using the slip knot technique in thoracoscopic anatomic segmentectomy. *J Vis Surg* 2017;3:100.